

CAPVT III

IONI

# Monoatomni ioni

- Gubitak ili primanje elektrona u neutralni atom
- Postoje samo u plinovitoj fazi

# Kationi u otopini


- Solvatacija
- Za neelektronegativne ione ovisna o naboju i radijusu – Latimer/hidratacija

$$\Delta H_{\text{hid}} \approx -60900 Z^2 / [(r + 50) / \text{pm}] \text{ kJ mol}^{-1}$$



- Ako je  $\chi > 1,5$  hidratacijska entalpija raste – kemijska veza između atoma i molekule otapala

# Akvatizirani kationi

- Molekule vode vezane ne centralni ion
- Raste kiselost (pada  $pK_a$ ) takvih molekula vode – “hidroliza”



(a)




$$K_a = 1,1 \cdot 10^{-5}$$



# Kategorije kiselosti iona

- $pK_a > 14$  nekiseli kationi ( $\text{Cs}^+$ ,  $\text{Rb}^+$ )
- $11,5 < pK_a < 14$  slabašno kiseli (*feebly acidic*;  $\text{Li}^+$ ,  $\text{Ba}^{2+}$ )
- $6 < pK_a < 11,5$  slabo kiseli ( $\text{Mg}^{2+}$ )
- $1 < pK_a < 6$  srednje kiseli ( $\text{Al}^{3+}$ )
- $-4 < pK_a < 1$  jako kiseli ( $\text{Ti}^{4+}$ )
- $pK_a < -4$  vrlo jako kiseli – kvantitativno reagiraju s vodom, postoje samo formalno ( $\text{Mn}^{7+}$ ,  $\text{Cr}^{6+}$ )



(a) Nonacidic Cations ( $pK_a > 14$ )





(b) Feebly Acidic Cations ( $pK_a = 11.5-14$ )





(c) Weakly Acidic Cations ( $pK_a = 6-11.5$ )



(d) Moderately Acidic Cations ( $pK_a = 1-6$ )



(e) Strongly Acidic Cations ( $-4 \leq pK_a \leq 1$ )




# Kiselost i svojstva iona

- Kiselost je funkcija naboja, radijusa (i elektronegativnosti, i drugih stvari koje se u prvu ruku mogu zanemariti...)

$$pK_a \approx 15,14 - 88,16 \text{ pm} * [Z^2/r]$$

za elektronegativnije elemente ( $\chi_P > 1,50$ ), potrebno je uvesti i korekciju na elektronegativnost:

$$pK_a \approx 15,14 - 88,16 \text{ pm} * [Z^2/r + 0,096 \text{ pm}^{-1} (\chi_P - 1,50)]$$



Korelacija mjerenih pK<sub>a</sub> i  
(korigiranog) omjera kvadrata  
naboja i radijusa

$[Z^2/r + 0,096 \text{ pm}^{-1} (\chi_P - 1,50)]$   
za monoatomne katione



# Anioni

- Također solvatirani
- Latimer/hidratacija  
$$\Delta H_{\text{hid}} \approx -57 \cdot 10^3 Z^2 / r \text{ pm kJ mol}^{-1}$$
- Ponašaju se kao baze

- $pK_b > 14$  nebazni anioni ( $\text{Cl}^-$ ,  $\text{Br}^-$ ,  $\text{I}^-$ )
- $11,5 < pK_b < 14$  slabašno bazni
- $6 < pK_b < 11,5$  slabo bazni ( $\text{F}^-$ )
- $1 < pK_a < 6$  srednje bazni ( $\text{Te}^{2-}$ )
- $-4 < pK_a < 1$  jako bazni ( $\text{Se}^{2-}$ ,  $\text{S}^{2-}$ )
- $pK_a < -4$  vrlo jako bazni ( $\text{O}^{2-}$ )







Figure 2.10

Acid–base predominance diagrams for (1 M total concentrations of) nonmetal monoatomic anions and their partially protonated and fully protonated forms.

- Baznost kao funkcija naboja i radijusa

$$pK_b \approx 29 - 1200 Z^2 / r$$





Figure 2.10

Acid–base predominance diagrams for (1 M total concentrations of) nonmetal monoatomic anions and their partially protonated and fully protonated forms.


# Što je kiselina?




Ono što je kiselo




Ono od čega ekstrakt  
ljubičice pocrveni



Ono što u vodenoj otopini  
poveća koncentraciju  
vodikovih kationa



Ono što otpušta  
protone (hidrone)



Ono što prima  
elektronski par

# Tvrde i meke kiseline i baze

- Za potpun opis ponašanja dane Lewisove kiseline nedovoljan je podatak o jakosti
- Pearson – tvrde i meke kiseline i baze
- Tvrde:
  - Malen radijus, velik naboj, mala polarizabilnost
- Meke:
  - Velik radijus, malen naboj, velika polarizabilnost

## Kiseline

tvrde	meke
H <sup>+</sup>	CH <sub>3</sub> Hg <sup>+</sup> , Hg <sup>2+</sup> , Hg <sub>2</sub> <sup>2+</sup>
Li <sup>+</sup> , Na <sup>+</sup> , K <sup>+</sup>	Pt <sup>2+</sup>
Ti <sup>4+</sup>	Pd <sup>2+</sup>
Cr <sup>3+</sup> , Cr <sup>6+</sup>	Ag <sup>+</sup>
BF <sub>3</sub>	BH <sub>3</sub>
R <sub>3</sub> C <sup>+</sup>	M <sup>0</sup>
	Au <sup>+</sup>

## Baze

tvrde	meke
OH <sup>-</sup>	H <sup>-</sup>
RO <sup>-</sup>	RS <sup>-</sup>
F <sup>-</sup> , Cl <sup>-</sup>	I <sup>-</sup>
NH <sub>3</sub>	PR <sub>3</sub>
CH <sub>3</sub> COO <sup>-</sup>	SCN <sup>-</sup>
CO <sub>3</sub> <sup>2-</sup>	CO
N <sub>2</sub> H <sub>4</sub>	C <sub>6</sub> H <sub>6</sub>

# Apsolutna tvrdoća

## Kiseline

## Baze

$$\eta = \frac{1}{2} \left[ \frac{\partial^2 E}{\partial^2 N} \right]_Z$$

$$\eta \approx -\frac{1}{2} (E_i + E_a) = -\frac{1}{2} \chi_M$$


H <sup>+</sup>	-	F <sup>-</sup>	7
Al <sup>3+</sup>	45.8	NH <sub>3</sub>	6.8
Li <sup>+</sup>	35.1	H <sup>-</sup>	6.8
Sc <sup>3+</sup>	24.6	CO	6.0
Na <sup>+</sup>	21.1	OH <sup>-</sup>	5.6
La <sup>3+</sup>	15.4	CN <sup>-</sup>	5.3
Zn <sup>2+</sup>	10.8	PH <sub>3</sub>	5.0
CO <sub>2</sub>	10.8	NO <sub>2</sub> <sup>-</sup>	4.5
SO <sub>2</sub>	5.6	SH <sup>-</sup>	4.1
I <sub>2</sub>	3.4	CH <sub>3</sub> <sup>-</sup>	4.0





# Višeatomni ioni

- Načelni pristup: centralni atom (“kation”) na koji su vezani drugi
- Koordinacijski broj – broj atoma vezanih na centralni atom
- Totalni koordinacijski broj – koordinacijski broj + broj elektronskih parova
- Maksimalni totalni koordinacijski broj – funkcija veličine centralnog atoma: 2 perioda – 4; 3. i 4. – 6; 5. i 6. – više od 6

# Fluoro-anioni






$\text{BF}_4^-$



$\text{IF}_7^-$

# Okso-anioni

- Totalni koordinacijski broj u pravilu za 1 manji od maksimalnog – elektronski a ne sterički razlozi



# klasifikacija

Classification	Type	Calculated $pK_{b1}$	Examples with Known $pK_{b1}$ Values
Nonbasic anions	$MO_4^-$	22.6	M = Cl, Br, Mn, Tc, Re
	$MO_3^-$	16.9	M = N, Cl, Br, I(13.2)
Feebly basic anions	$MO_4^{2-}$	12.4	M = S(12.1), Se(12.0), Xe, Cr(7.5), Mo(9.9), W(9.4), Fe(6.2), Ru, Os
	$MO_2^-$	11.2	M = N(10.7), Cl(12.1)
Moderately basic anions	$MO_6^{4-}$	3.4	M = Xe, Os
	$MO_4^{3-}$	2.2	M = P(2.0), As(1.5), V(1.0)
	$MO_3^{2-}$	6.7	M = C, S(6.8), Se(7.4), Te(6.3)
	$MO^-$	5.5	M = Cl(6.5), Br(5.3), I(3.4)
Very strongly basic anions (Exist as hydroxo anions in solution)	$MO_6^{5-}$	-6.8	M = I, Np
	$MO_4^{4-}$	-8.0	M = Si, Ge
	$MO_3^{3-}$	-3.5	M = As, Sb
	$MO_6^{6-}$	-17.0	M = Te
	$MO_4^{5-}$	-18.2	M = B, Al, Ga
	$MO_3^{4-}$	-14.7	M = Sn

SOURCES: Known  $pK_{b1}$  values are calculated from the appropriate  $pK_a$  values given in F. A. Cotton and G. Wilkinson, *Advanced Inorganic Chemistry: A Comprehensive Text*, 5th ed., Wiley-Interscience, New York; 1988, p. 105; R. C. Weast, Ed., *Handbook of Physics and Chemistry*, 50th ed., Chemical Rubber Publishing Co., Cleveland, 1969; J. A. Dean, Ed., *Lange's Handbook of Chemistry*, 13th ed., New York, McGraw-Hill, 1985; and B. H. J. Bielski, *Free Radical Res. Commun.*, **12-13**, 469 (1991).


# Baznost okso-aniona

- Raste s nabojem ( $pK_b$  pada za oko 10,2 za svaki dodatni negativni naboj)
- Pada s brojem kisikovih atoma ( $pK_b$  raste za oko 5,7 za svaki dodatni kisikov atom)
- Pada s porastom elektronegativnosti centralnog atoma

za ion  $MO_x^{y-}$

$$pK_b = 10 + 5,7x - 10,2y \pm 1,0$$

# Protoniranje okso-aniona



$$\Delta pK_b = -5,7 + 10,2 = 4,5$$

Nastavljajući se do kiseline  $[\text{MO}_{(x-y)}(\text{OH})_y]$

$$pK_a = 14 - pK_b = 8,5 - 5,7(x-y)$$

$x - y$  = broj okso-liganada u kiselinu - nužno poznavanje strukture ( $\text{H}_3\text{PO}_3$  i  $\text{H}_3\text{PO}_2$ ;  $pK_a \approx 2$ )