

Biomarker pigment divinyl chlorophyll *a* as a tracer of water masses?

Maja Mejdandžić¹, Hrvoje Mihanović², Tina Šilović³, Jorijntje Henderiks⁴, Luka Šupraha⁵, Dorotea Polović¹, Sunčica Bosak¹, Ivana Bošnjak¹, Ivona Cetinić⁶, Goran Olujić⁷, Zrinka Ljubešić¹

¹University of Zagreb, Faculty of Science, Department of Biology, Rooseveltov trg 6, 10000 Zagreb, Croatia (zrinka.ljubesic@biol.pmf.hr)
²Institute for Oceanography and Fisheries, Setalište I. Meštrovića 63, 21000 Split, Croatia
³Center for Marine Research, Ruđer Bošković Institute, G. Paliaga 5, 52210 Rovinj, Croatia
⁴CEES, Dept. of Biosciences, University of Oslo, P.O. Box 1066 Blindern, 0316 Oslo, Norway
⁵Department of Earth Sciences, Paleobiology Programme, Uppsala University, Villavägen 16, SE-752 36 Uppsala, Sweden.
⁶NASA, 300 E St SW, Washington, DC 20546, USA

The ecological preferences of different phytoplankton types drive their temporal and spatial distributions, reflecting their dependence on certain temperature ranges, light levels, nutrient availability and other environmental gradients. Hence, some phytoplankton taxa can be used as water mass tracers (biotracers).

In order to determine the biotracers of targeted Adriatic water masses, eight sampling campaigns have been conducted in the southern Adriatic (Albanian shelf, May 2009) and the middle Adriatic (November 2011, February, March, May and August 2012, February and July 2013) (Fig. 1.). The surveyed area is greatly influenced by the Levantine Intermediate Water (LIW) and East Adriatic Current (EAC). The warm and saline LIW represents a part of the EAC and enters the Adriatic from the Ionian Sea spreading northwards at the intermediate level, usually at depths between 100 and 600 m, with the core from 200 to 400 m. The spatial and temporal distribution, as well as the phytoplankton community composition was investigated in relation to the encountered environmental parameters (temperature (TEMP), salinity (SAL), density (Sigma T), nitrates (NO₃), nitrites (NO₂), ammonia (NH₄⁺), silicates (SiO₄⁻) and phosphates (PO₄³⁻). A combination of several taxonomical techniques; flow cytometry (FC) and high-performance liquid chromatography (HPLC) pigment analysis, allowed us to investigate phytoplankton community composition across all size fractions.




Fig. 1. Oceanographic surveys and sampling points. 1 - Albanian shelf, May 2009; 2 - Middle Adriatic, February and July 2013; 3 - Telašćica bay, middle Adriatic, October 2011, February, March, May and August 2012.




Fig. 2. Albanian shelf, May 2009. A. PCA analysis of physico-chemical parameters overlaid with pigment biomarker *dvchl a* concentration. *Dvchl a* was detected within the layer of LIW intrusion which is indicated by higher salinity values. B. *Dvchl a* as a good biomarker of *Prochlorococcus* presence in the water column was confirmed by flow-cytometry analysis.




Fig. 3. Middle Adriatic, Stations AD1, AD2, AD3, February and July 2013. PCA analysis of physico-chemical parameters overlaid with pigment biomarker *dvchl a* concentration. *Dvchl a* was detected within the layer of LIW intrusion which is indicated by higher salinity and density values.




Fig. 4. Telašćica bay, Middle Adriatic 2011/2012. A. PCA analysis of physico-chemical parameters. Temperature (seasons) was recognized as main driving force of this ecological system which is shown with separation of samples into three groups: I - winter and spring, II - summer and autumn and III - spring and summer. Different seasons shown in separate PCA analysis (WINTER, AUTUMN, SPRING and SUMMER). PCA analysis of physico-chemical parameters overlaid with pigment biomarker *dvchl a* concentration confirmed *dvchl a* signal within the layer of LIW intrusion and its correlation with higher salinity and density values. In winter conditions it is suggested that LIW intrusion was not as pronounced in middle Adriatic and *dvchl a* signal was not correlated with density.

The overall phytoplankton spatial distribution demonstrated high patchiness. However, in certain seasons, divinyl chlorophyll *a* (*dvchl a*), a pigment specific for the prokaryote *Prochlorococcus*, was found in high concentrations below 50 m – the portion of the water column in which LIW was frequently detected. The presence of *Prochlorococcus* was confirmed by flow cytometry, corroborating its important role in the formation of the deep chlorophyll maximum (Fig. 2 - 4.).

Based on these results, we hypothesize that the picophytoplanktonic prokaryote *Prochlorococcus*, that is easily detectable in the water column owing to its specific pigment structure, can be used for tracing Levantine Intermediate Water in the Adriatic.

ACKNOWLEDGMENTS

This work has been supported in part by Croatian Science Foundation under the project 6433, Research Council of Norway (FRIMEDBIO prijet 197823) and Royal Swedish Academy of Sciences through a grant from the Knut and Alice Wallenberg Foundation (KAW 2009.0287) and Norwegian Cooperation Programme on Research and Higher Education with countries in the Western Balkans: 'Marine Science and Coastal Management in the Adriatic, Western Balkans'. We thank to PP 'Telašćica' and Hydrographic Institute of the Republic of Croatia for their help during the fieldwork. M.M. has been supported by University of Zagreb staff exchange mobility grant.